

University of Groningen

The microenvironment of Hodgkin lymphoma

Sattarzadeh, Ahmad

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Sattarzadeh, A. (2016). *The microenvironment of Hodgkin lymphoma: Composition and interaction*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Stellingen

Behorende bij het proefschrift

The microenvironment of Hodgkin Lymphoma

Composition and interaction

1. The interaction between the neoplastic cells and the cells of the microenvironment in Hodgkin lymphoma is a matter of survival or death (This thesis)
2. High numbers of CD8+ T-cells and NK cells in classical Hodgkin lymphoma indicates an Epstein-Barr virus positive tumor (This thesis)
3. Being within or outside of the tumor cell area. That should be the question (This thesis)
4. Increase in the number of follicular T helper cells in the microenvironment of nodular lymphocyte predominant Hodgkin lymphoma is caused by the absence of regulatory T-cells (This thesis)
5. Lymphocyte-predominant cells are protected by presence of rosetting CD57+ and CD57-PD-1+ TFH cells (This thesis)
6. The microenvironments of classical and nodular lymphocyte predominant Hodgkin lymphoma have both similarities and differences (This thesis)

Ahmad Sattarzadeh, 12 September 2016